Public reporting burden for the collection of information maintaining the data needed, and completing and review including suggestions for reducing this burden, to Washi VA 22202-4302 Respondents should be aware that notw	ing the collect ngton Headqu	ion of information Send comments arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 01 JAN 2005	2. REPORT TYPE <b>N/A</b>		3. DATES COVERED		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Effectiveness of local antibiotic delivery with osteoinductive osteoconductive bone-graft substitutes.				5b. GRANT NUMBER	
6. AUTHOR(S)  Beardmore A. A., Brooks D. E., Wenke J. C., Thomas D. B.,				5c. PROGRAM ELEMENT NUMBER	
				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  United States Army Institute of Surgical Research, JBSA Fort Sam houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATE Approved for public release, di		on unlimited			
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF: 17. LIN				18. NUMBER	19a. NAME OF
a REPORT b ABSTRA unclassified unclassi		c THIS PAGE unclassified	- ABSTRACT SAR	OF PAGES  1	RESPONSIBLE PERSON

**Report Documentation Page** 

Form Approved OMB No. 0704-0188 J Bone Joint Surg Am. 2005 Jan;87(1):107-12.

## Effectiveness of local antibiotic delivery with an osteoinductive and osteoconductive bone-graft substitute

Beardmore AA, Brooks DE, Wenke JC, Thomas DB.

United States Army Institute of Surgical Research, Fort Sam, Houston, Texas, USA. anthony.beardmore@cen.amedd.army.mil

BACKGROUND: The morbidity associated with open fractures and open fracture treatment is well established. An osteoinductive and osteoconductive bone-graft substitute that prevents infection would decrease the number of procedures required to treat contaminated fractures by eliminating the need for surgical removal of cement beads and perhaps autograft harvest. We hypothesized that the combination of tobramycin-impregnated calcium sulfate pellets and demineralized bone matrix would prevent the establishment of infection in a contaminated fracture model. METHODS: A unicortical 12-mm-diameter defect was created in the proximal tibial metaphysis of twenty-nine Spanish goats. After contaminating the wounds with an infective dose of Staphylococcus aureus, we divided the animals into four groups. The negative control group received no treatment, the positive control group received tobramycin-impregnated polymethylmethacrylate beads, the demineralized bone matrix group received 2.5 mL of demineralized bone matrix, and the experimental group received tobramycin-impregnated calcium sulfate pellets with 2.5 mL of demineralized bone matrix. Radiographs were made and intraosseous tissue cultures were performed on postoperative day 21. RESULTS: The cultures showed no evidence of intramedullary infection in the experimental or the positive control group, but they were positive for Staphylococcus aureus in six of the seven goats in the negative control group and seven of the eight goats in the demineralized bone matrix group. CONCLUSIONS: The combination of tobramycin-impregnated calcium sulfate pellets and demineralized bone matrix was effective in preventing intramedullary Staphylococcus aureus infection in a contaminated goat fracture model.